



# City of Milford, Connecticut

- Founded 1639 -

## HEALTH DEPARTMENT

A. Dennis McBride, M.D., M.P.H.  
Health Director

Telephone: (203) 783-3285  
FAX: (203) 783-3286  
E-mail: admcbride@ci.milford.ct.us

January 6, 2011

Dear Parents,

The Milford Health Department and Milford Public Schools are currently working together closely to investigate an outbreak of gastrointestinal illness in the school setting. The symptoms of the illness are consistent with Norwalk-like viruses, or Norovirus. The most common symptoms of Norovirus infection include nausea, vomiting, diarrhea, and abdominal cramps. Other symptoms can include headache, fever, chills, and muscle aches. Most people get better within 1-2 days. The primary mode of transmission is fecal-oral spread through person-to-person contact. Similar symptoms are currently being seen in various settings across the state and are also presumed to be Norovirus.

While Norovirus is not usually considered a life-threatening illness, symptoms can be debilitating to the extent that individuals will not be able to attend work or school. As such, the Milford Health Department and Milford Public Schools are taking the following measures to control the spread of this virus across the school district. Please note that these measures will not eliminate the virus, however the actions taken will serve to reduce the transmission of the virus from person to person.

- **Milford Health Department staff has conducted site visits of all Milford Public Schools to ensure that all food preparation and handling areas are safe.** All food workers have been instructed to stay at home if they experience any symptoms and/or have household contacts that are ill.
- **All Milford Public Schools will be thoroughly cleaned and disinfected using appropriate cleaners.** Any and all potentially contaminated surfaces in each facility throughout the district will be properly cleaned and disinfected immediately.
- **All students and staff will be required to wash their hands with soap and water for 20 seconds upon arrival at school, as well as before and after lunch.** There is no specific treatment or vaccine to prevent Norovirus infection. The most effective preventive measure is to practice proper hand hygiene by washing hands with soap and warm water for 20 seconds, especially after using the toilet and before handling food.
- **Students and staff who are sick should remain at home.** Sick individuals should avoid contact with others to decrease the spread of this virus. If your child comes to school with symptoms of this virus, he/she will be sent home.

Attached is some basic information regarding Norovirus and its transmission. The above measures have been taken to control the spread of this virus. No further measures are required at this time. The Milford Health Department and Milford Public Schools will continue to closely monitor this situation and update parents as necessary.

Thank you for helping us keep our schools as healthy as possible for all students and staff.

Sincerely,

A. Dennis McBride, MD, MPH  
Health Director

Michael Cummings  
Acting Superintendent of Schools

# Norovirus Illness: Key Facts



## Norovirus—the stomach bug

- Norovirus is a highly contagious illness caused by infection with a virus called norovirus. It is often called by other names, such as viral gastroenteritis, stomach flu, and food poisoning.
- Norovirus infection causes acute gastroenteritis (inflammation of the stomach and intestines); the most common symptoms are diarrhea, vomiting, and stomach pain.
- Anyone can get norovirus, and they can have the illness multiple times during their lifetime.
- Norovirus is the most common cause of acute gastroenteritis in the United States.

## Norovirus illness can be serious

- Norovirus can make people feel extremely ill and vomit or have diarrhea many times a day.
- Most people get better within 1 to 2 days.
- Dehydration can be a problem among some people with norovirus infection, especially the very young, the elderly, and people with other illnesses.

## Norovirus is highly contagious and spreads rapidly

- Noroviruses are highly contagious, and outbreaks are common due to the ease of transmission.
- People with norovirus are contagious from the moment they begin feeling ill to at least 3 days and perhaps for as long as 2 weeks after recovery, making control of this disease even more difficult.
- Norovirus can spread rapidly in closed environments like daycare centers and nursing homes.

## Many sources for norovirus infection

Noroviruses are found in the stool and vomit of infected people. People can become infected by

- Eating food or drinking liquids that are contaminated with norovirus.
- Touching surfaces or objects that are contaminated with norovirus, and then placing their hand in their mouth.
- Having direct contact with an infected person; for example, by exposure to the virus when caring for or when sharing food, drinks, or eating utensils with an infected person.

National Center for Immunization and Respiratory Diseases  
Division of Viral Diseases





## Tips to prevent the spread of norovirus

**Practice proper hand hygiene:** Wash your hands carefully with soap and water, especially after using the toilet and changing diapers and always before eating or preparing food. Alcohol-based hand sanitizers (containing at least 62% ethanol) may be a helpful addition to hand washing, but they are not a substitute for washing with soap and water.

See Handwashing: Clean Hands Save Lives (<http://www.cdc.gov/cleanhands/>) for more information about washing hands.

**Take care in the kitchen:** Carefully wash fruits and vegetables, and cook oysters and other shellfish thoroughly before eating them.

**Do not prepare food while infected:**

People who are infected with norovirus should not prepare food for others while they have symptoms and for 3 days after they recover from their illness.

**Clean and disinfect contaminated surfaces:** After an episode of illness, such as vomiting or diarrhea, immediately clean and disinfect contaminated surfaces by using a bleach-based household cleaner as directed on the product label or a solution made by adding 5–25 tablespoons of household bleach to 1 gallon of water.

**Wash laundry thoroughly:** Immediately remove and wash clothing or linens that may be contaminated with vomit or fecal matter. Handle soiled items carefully—without agitating them—to avoid spreading virus. They should be laundered with detergent at the maximum available cycle length and then machine dried.

## No specific treatment or vaccine for norovirus infection

- There is no vaccine to prevent norovirus infection, although this is an area of active research.
- There is no specific drug to treat people with norovirus illness.
- Rehydration is important for infected people—they must drink plenty of liquids to replace fluid lost through vomiting and diarrhea. In some cases, fluid may need to be given intravenously.

